

ON THE COMPLEMENTARITY PRINCIPLE AND THE UNCERTAINTY PRINCIPLE

E. S. Guerra

Departamento de Física

Universidade Federal Rural do Rio de Janeiro

Cx. Postal 23851, 23890-000 Seropédica, RJ, Brazil

email: emerson@ufrj.br

February 1, 2008

Abstract

We present a scheme in which we investigate the two-slit experiment and we show that the principle of complementarity is more fundamental than the uncertainty principle.

PACS: 03.65.Ud; 03.67.Mn; 32.80.-t; 42.50.-p

Keywords: principle of complementarity, uncertainty principle, cavity QED.

The investigation of the double slit experiment in which we have two micromaser cavities, each one associated with one of the slits, was proposed before by Scully and Walther in a series of very interesting articles where they investigate the principle of complementarity and the uncertainty principle [1, 2] and they conclude that the principle of complementarity is more fundamental than the uncertainty principle. The same conclusion can be reached very easily using the scheme presented below. We are going to consider a screen with two slits $SL1$ (at ζ_1) and $SL2$ (at ζ_2) with two cavities $C1$ and $C2$ behind respectively each slit and prepared respectively in an even coherent state $|+\rangle_1$ and an odd coherent state $|-\rangle_2$, where

$$|\pm\rangle_k = |\alpha_k\rangle \pm |-\alpha_k\rangle \quad (1)$$

[3], through which fly Rydberg atoms of relatively long radiative lifetimes [4]. We also assume perfect microwave cavities, that is, we neglect effects due to decoherence. Concerning this point, it is worth to mention that nowadays it is possible to build up niobium superconducting cavities with high quality factors Q . It is possible to construct cavities with quality factors $Q \sim 10^8$ [5]. Even cavities with quality factors as high as $Q \sim 10^{12}$ have been reported [6], which, for frequencies $\nu \sim 50$ GHz gives us a cavity field lifetime of the order of a few seconds.

Let us first show a possible way of preparing the cavities in the states (1). Suppose we prepare cavity Ck initially in a coherent state $|i\alpha_k\rangle$. Then, we prepare a two-level atom $A0$ in a coherent superposition, preparing it initially in state $|f_0\rangle$, passing it through a first Ramsey zone $R1$ where

$$R_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} c_f & c_e \\ -c_e & c_f \end{bmatrix}, \quad (2)$$

and we get

$$|\psi\rangle_0 = c_e |e_0\rangle + c_f |f_0\rangle. \quad (3)$$

After that, $A0$ flies through cavity Ck . The $|e_0\rangle \rightleftharpoons |f_0\rangle$ transition is far from resonance with the cavity central frequency such that only virtual transitions occur between these states and the interaction of the atom with the cavity mode in $C1$ is described by the time evolution operator [7].

$$U(t) = e^{-i\varphi(a^\dagger a + 1)} |e\rangle\langle e| + e^{i\varphi a^\dagger a} |f\rangle\langle f|, \quad (4)$$

where a ($a^\dagger$) is the annihilation (creation) operator, $\varphi = g^2\tau/\Delta$, g is the coupling constant, $\Delta = \omega_e - \omega_f - \omega$ is the detuning where ω_e and ω_f are the frequency of the upper and lower levels respectively and ω is the cavity field frequency and τ is the atom-field interaction time. After the atom passes through $C1$ the state of the system $A0 - C1$, for $\varphi = \pi/2$, is given by

$$|\psi\rangle_{A0-C1} = -ic_e |e_0\rangle |\alpha_k\rangle + c_f |f_0\rangle |-\alpha_k\rangle.$$

Then, the atom enters a second Ramsey zone $R2$ where we have

$$R_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -i \\ i & 1 \end{bmatrix}, \quad (5)$$

that is,

$$\begin{aligned} |e_0\rangle &\rightarrow \frac{1}{\sqrt{2}}(|e_0\rangle + i|f_0\rangle), \\ |f_0\rangle &\rightarrow \frac{1}{\sqrt{2}}(-i|e_0\rangle + |f_0\rangle). \end{aligned} \quad (6)$$

After the atom cross the Ramsey zone $R2$, the state of the system $A0 + Ck$ is given by

$$\begin{aligned} |\psi\rangle_{A0-Ck} &= \frac{1}{\sqrt{2}}[-ic_e |\alpha_k\rangle - ic_f |-\alpha_k\rangle] |e_0\rangle \\ &\quad + \frac{1}{\sqrt{2}}[c_e |\alpha_k\rangle + c_f |-\alpha_k\rangle] |f_0\rangle, \end{aligned}$$

and for $c_e = c_f$ we get $|\psi\rangle_{Ck} = |+\rangle_k$ if we detect $|e_0\rangle$ or $|f_0\rangle$ and for $c_e = -c_f$ we get $|\psi\rangle_{Ck} = |-\rangle_k$ if we detect $|e_0\rangle$ or $|f_0\rangle$.

Now, for a three-level lambda atom interacting with the electromagnetic field inside a cavity where the upper and the two degenerated lower states are $|a\rangle$, $|b\rangle$ and $|c\rangle$ respectively, and for which the $|a\rangle \rightleftharpoons |c\rangle$ and $|a\rangle \rightleftharpoons |b\rangle$ transitions are in the far from resonance interaction limit, the time evolution operator $U(t)$ for the atom-field interaction in a cavity Ck is given by [8]

$$\begin{aligned} U(\tau) &= \frac{1}{2}(e^{i\varphi a_k^\dagger a_k} + 1)|b\rangle\langle b| + \frac{1}{2}(e^{i\varphi a_k^\dagger a_k} - 1)|b\rangle\langle c| + \\ &\quad \frac{1}{2}(e^{i\varphi a_k^\dagger a_k} - 1)|c\rangle\langle b| + \frac{1}{2}(e^{i\varphi a_k^\dagger a_k} + 1)|c\rangle\langle c|. \end{aligned} \quad (7)$$

where a_k ($a_k^\dagger$) is the annihilation (creation) operator for the field in cavity Ck , $\varphi = 2g^2\tau/\Delta$, g is the coupling constant, $\Delta = \omega_a - \omega_b - \omega = \omega_a - \omega_c - \omega$ is the detuning where ω_a , ω_b and ω_c are the frequency of the upper and of the two degenerate lower levels respectively and ω is the cavity field frequency and τ is the atom-field interaction time. For $\varphi = \pi$, we get

$$U(\tau) = -\exp(i\pi a_k^\dagger a_k) |a\rangle\langle a| + \Pi_{k,+}|b\rangle\langle b| + \Pi_{k,-}|b\rangle\langle c| + \Pi_{k,-}|c\rangle\langle b| + \Pi_{k,+}|c\rangle\langle c|, \quad (8)$$

where

$$\begin{aligned}\Pi_{k,+} &= \frac{1}{2}(e^{i\pi a_k^\dagger a_k} + 1), \\ \Pi_{k,-} &= \frac{1}{2}(e^{i\pi a_k^\dagger a_k} - 1),\end{aligned}\tag{9}$$

and we have

$$\begin{aligned}\Pi_{k,+}|+\rangle_k &= |+\rangle_k, \\ \Pi_{k,+}|-\rangle_k &= 0, \\ \Pi_{k,-}|-\rangle_k &= -|-\rangle_k, \\ \Pi_{k,-}|+\rangle_k &= 0,\end{aligned}\tag{10}$$

which are easily obtained from Eqs. (9) and (1) using $e^{za_k^\dagger a_k}|\alpha_k\rangle = |e^z\alpha_k\rangle$ [9]. Let us assume that we send three-level lambda atoms through the slits and cavities. Consider an atom $A1$ prepared in the state $|b_1\rangle$ flying through the double slit. Before $A1$ crosses the cavities we have

$$|\psi\rangle_{A1-SL1-SL2} = \frac{1}{\sqrt{2}}(|\zeta_1\rangle + |\zeta_2\rangle)|+\rangle_1|-\rangle_2 |b_1\rangle\tag{11}$$

and after it has interacted with $C1$ and $C2$, taking into account (8),

$$|\psi(t_0)\rangle_{A1-C1-C2} = \frac{1}{\sqrt{2}}(|\zeta_1\rangle |b_1\rangle - |\zeta_2\rangle |c_1\rangle)|+\rangle_1|-\rangle_2.\tag{12}$$

Now, writing $\psi(x, t)_{A1-C1-C2} = \langle x|U(t, t_0)|\psi(t_0)\rangle_{A1-C1-C2}$, $\psi_1(x, t) = \langle x|U(t, t_0)|\zeta_1\rangle$ and $\psi_2(x, t) = \langle x|U(t, t_0)|\zeta_2\rangle$, where $|x\rangle$ is a point on a screen in front of the double slit screen at a certain distance L from it, we have

$$\psi(x, t)_{A1} = \frac{1}{\sqrt{2}}\{\psi_1(x, t) |b_1\rangle - \psi_2(x, t) |c_1\rangle\}.\tag{13}$$

and

$$|\psi(x, t)_{A1}|^2 = \frac{1}{2}[|\psi_1(x, t)|^2 + |\psi_2(x, t)|^2]\tag{14}$$

since $\langle c_1 | b_1\rangle = 0$ and if there were no cavities we would obtain

$$|\psi(x, t)_{A1}|^2 = \frac{1}{2}[|\psi_1(x, t)|^2 + |\psi_2(x, t)|^2 + 2Re\{\psi_1^*(x, t)\psi_2(x, t)\}]\tag{15}$$

which presents interference fringes. Therefore, when we place cavities $C1$ and $C2$ prepared in the states $|+\rangle_1$ and $|-\rangle_2$ respectively, the interference fringes are washed out. This happens because the parity information of the cavities is transferred to the internal state of the atom. Notice that if we detect the atomic state of $A1$ after it has crossed the slits and before it strikes the detection screen at $|x\rangle$ and we find $|b_1\rangle$, we can say that the atom has passed through slit $SL1$, and if we detect $|c_1\rangle$, we can say that the atom has passed through slit $SL2$ and we get which-path information (particle behavior) detecting the atomic state. That is, assuming that the detection of the internal states does not disturb the external state of motion of the centre of mass of the atom, in the case we detect $|b_1\rangle$ we get

$$|\psi(x, t)_{A1}|^2 = |\psi_1(x, t)|^2,\tag{16}$$

and in the case we detect $|c_1\rangle$ we get

$$|\psi(x, t)_{A1}|^2 = |\psi_2(x, t)|^2. \quad (17)$$

Therefore, the cavities allow us to get which-path information. Notice that as the atom-field interaction is dispersive, we can state that the uncertainty principle plays no role at all. Therefore, we get the same conclusion of Scully and Walther [1], that is, the complementarity principle is more fundamental than the uncertainty principle since as the experiment cited and the present experiment represent a way around the uncertainty principle.

Now, assume that we place a Ramsey cavity on the way of the atom just before it passes through the slits and cavities so that the atomic state before the atom crosses the cavities be

$$|\psi\rangle_{A1} = \frac{1}{\sqrt{2}}(|b_1\rangle + |c_1\rangle). \quad (18)$$

In this case the initial state of the system is

$$|\psi\rangle_{A1-SL1-SL2} = \frac{1}{2}(|\zeta_1\rangle + |\zeta_2\rangle)(|+\rangle_1|-\rangle_2(|b_1\rangle + |c_1\rangle)) \quad (19)$$

and after the atom crosses the cavities

$$|\psi\rangle_{A1-C1-C2} = \frac{1}{2}[|\zeta_1\rangle(|b_1\rangle + |c_1\rangle) - |\zeta_2\rangle(|c_1\rangle + |b_1\rangle)](|+\rangle_1|-\rangle_2) \quad (20)$$

and therefore we have

$$|\psi(x, t)_{A1}|^2 = \frac{1}{4}\{|\psi_1(x, t)|^2 + |\psi_2(x, t)|^2 - 2\text{Re}\{\psi_1^*(x, t)\psi_2(x, t)\}\} \quad (21)$$

and we have interference as we would expect since the atom enters the cavities in an atomic state superposition of their internal states and no information about the state of the cavities can be transferred to the atom which exits the cavities also in an atomic state superposition which do not allows us to get which-path information detecting the atomic state. Note that, in the case we do not have a Ramsey cavity and we prepare the atom, for instance, in state $|b_1\rangle$, we know that it is in this state and, if we accept the theoretical formalism of quantum mechanics as correct, according to it we know that in this case, as we have seen above, the parity information of the cavities is transferred to the atom allowing us to get which-path information.

Let us consider now a scheme of atomic states similar the one used above to prepare the cavities in an even or an odd coherent state, but now we take a three-level cascade atom with $|e\rangle$, $|f\rangle$ and $|g\rangle$ being the upper, intermediate and lower atomic states. Again we assume that the transition $|f\rangle \rightleftharpoons |e\rangle$ is far enough from resonance with the cavity central frequency such that only virtual transitions occur between these states. In addition we assume that the transition $|e\rangle \rightleftharpoons |g\rangle$ is highly detuned from the cavity frequency so that there will be no coupling with the cavity field (only the states $|f\rangle$ and $|e\rangle$ interact with the field in the cavity). Here we are going to consider the effect of the atom-field interaction taking into account only levels $|f\rangle$ and $|g\rangle$. We do not consider level $|e\rangle$ since it will not play any role in our scheme. Therefore, we have effectively a two-level system involving states $|f\rangle$ and $|g\rangle$. Considering levels $|f\rangle$ and $|g\rangle$ and taking into account (4), we can write an effective time evolution operator

$$U_k(t) = e^{i\varphi a^\dagger a} |f\rangle\langle f| + |g\rangle\langle g|, \quad (22)$$

where the second term above was put by hand just in order to take into account the effect of level $|g\rangle$. Let us assume that atom $A1$ is prepared in state $|g\rangle$ and, on the way to the screen with two slits and two cavities, there is a Ramsey cavity $R1$ where the atomic states are rotated according to

$$R_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (23)$$

that is, after $A1$ crosses this Ramsey cavity we have

$$|\psi\rangle_{A1} = \frac{1}{\sqrt{2}}(|f_1\rangle + |g_1\rangle), \quad (24)$$

and, for $\varphi = \pi$, after $A1$ flies through slit $SL1$ and cavity $C1$ and through slit $SL2$ and cavity $C2$, we have

$$|\psi(t_0)\rangle_{A1-C1-C2} = \frac{1}{2}\{|\zeta_1\rangle(|f_1\rangle + |g_1\rangle) + |\zeta_2\rangle(-|f_1\rangle + |g_1\rangle)\}|+\rangle_1|-\rangle_2, \quad (25)$$

and taking into account the time evolution operator $U(t, t_0)$

$$|\psi_1(x, t)_{A1}|^2 = \frac{1}{4}[|\psi_1(x, t)|^2 + |\psi_2(x, t)|^2], \quad (26)$$

and we have no interference fringes since $(-\langle f_1| + \langle g_1|)(|f_1\rangle + |g_1\rangle) = 0$. Now, if we do not have $R1$ on the way of the atom to the two-slit screen we get

$$|\psi(t_0)\rangle_{A1-C1-C2} = \frac{1}{2}\{|\zeta_1\rangle|g_1\rangle - |\zeta_2\rangle|f_1\rangle\}|+\rangle_1|-\rangle_2 \quad (27)$$

and therefore,

$$|\psi(x, t)_{A1}|^2 = \frac{1}{2}[|\psi_1(x, t)|^2 + |\psi_2(x, t)|^2 - 2\text{Re}\{\psi_1^*(x, t)\psi_2(x, t)\}], \quad (28)$$

and we have interference fringes. In the case we have the Ramsey cavity on the way of the atom to the two-slit screen the atom enters the cavities $C1$ and $C2$ in an atomic state superposition and now the cavities transfer their parity information to the atom. We can see that we can get which-path information in this case if we consider another Ramsey cavity $R2$ where

$$R_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}, \quad (29)$$

and

$$\frac{1}{\sqrt{2}}(|f_1\rangle + |g_1\rangle) \longrightarrow |f_1\rangle, \quad (30)$$

$$\frac{1}{\sqrt{2}}(-|f_1\rangle + |g_1\rangle) \longrightarrow |g_1\rangle \quad (31)$$

just behind $C1$ or $C2$. If $R2$ is just after cavity $C1$ we can detect atom $A1$ in state $|f_1\rangle$ and we know that the atom has passed through slit $SL1$ and if $R2$ is just after cavity $C2$ we can detect atom $A1$ in state $|g_1\rangle$ and we know that the atom has passed through slit $SL2$, that is, we get which-path information. In the case we do not have $R1$ on the way of the atom to the slits, it is clear that we cannot get which-path information.

We should point out that the field of the cavities is not disturbed in any way in the above schemes and therefore, these experiments could be performed with any number of atoms which would allow us to get experimentally a two peaks pattern (particle behavior) or a wiggly pattern (wave behavior). Therefore, aside technical difficulties, these experiments could not be considered *gedanken* experiments. A technical problem in these experiments is related to the fact that the separation between the slits should be very small and one should work also with very small cavities. Although this is not an easy problem to deal with, we think that the schemes we have discussed are at least of academic interest. We should point out also that many experiments which were considered as *gedanken* experiments in the past turned out to be realized in laboratory nowadays.

We intend to publish a further investigation along this line elsewhere.

References

- [1] M. O. Scully and H. Walther, Phys. Rev. A **39**, 5229 (1989); M. O. Scully, B-G Englert and H. Walther, Nature **351**, 111 (1991); M. O. Scully and H. Walther, Phys. Rev. A **49**, 1562 (1994); M. O. Scully and H. Walther, Found. Phys. **28**, 399 (1998).
- [2] J. Baggott, New Sci. **133**, 36 (1992).
- [3] E. S. Guerra, B. M. Garraway and P. L. Knight, Phys. Rev. A **55**, 3482 (1997).
- [4] A. A. Radzig and B. M. Smirnov, *Reference Data on Atoms, Molecules, and Ions* (Springer-Verlag, Berlin, 1985); T. F. Gallagher, *Rydberg Atoms* (Cambridge Univ. Press, Cambridge, 1984).
- [5] M. Brune, F. Schmidt-Kaler, A. Maali, J. Dreyer, E. Hagley, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **76**, 1800 (1996).
- [6] G. Rempe, F. Schmidt-Kaler and H. Walther, Phys. Rev. Lett. **64**, 2783 (1990).
- [7] M. Orszag, *Quantum Optics*, (Springer-Verlag, Berlin, 2000).
- [8] P. L. Knight, Phys. Scr. **T12**, 51 (1986); S. J. D. Phoenix and P. L. Knight, J. Opt. Soc. Am. B **7**, 116 (1990).
- [9] W. H. Louisell, *Quantum Statistical Properties of Radiation*, (Wiley, New York, 1973).